



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$Cxy - Fx - Gy + H = 0 \dots\dots(7),$$

in which  $A=bc-f^2$ ,  $B=ca-g^2$ ,  $C=ab-h^2$ ,  $F=gh-af$ ,  $G=hf-bg$ ,  $H=fg-ch$ .

For the conic (4),  $A=-1/b^2$ ,  $B=-[(1+\lambda)/a^2]$ ,  $C=[(1-\lambda)/(a^2b^2)]$ ,  $F=-\lambda/a^2b^2$ ,  $G=H=0$ , and these in (6) and (7) give

$$(1-\lambda)(x^2-y^2)-2b\lambda y+b^2(1-\lambda)-a^2=0\dots\dots(8),$$

$$(1-\lambda)y+b\lambda=0\dots\dots(9).$$

Eliminating  $\lambda$  from (8) and (9), the required locus is

$$x^2+y^2+\frac{a^2-2b^2}{b}y=a^2-b^2\dots\dots(10).$$

This is a circle, radius  $a^2/2b$ , or one-half the radius of curvature of (1) at the extremity of the minor axis.

115. Proposed by MARY M. BLAINE, B. Sc., Graduate Student, Drury College, Springfield, Mo.

The locus of a point such that the sum of the squares of its normals form a given ellipsoid is constant, is a co-axial ellipsoid. [From *C. Smith's Solid Analytical Geometry*, page 95.]

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Let  $(l, m, n)$  be the point whose locus is required. Then from the conditions of the problem we have

$$\begin{aligned} &(l-x_1)^2+(l-x_2)^2+(l-x_3)^2+(l-x_4)^2+(l-x_5)^2+(l-x_6)^2 \\ &+(m-y_1)^2+(m-y_2)^2+(m-y_3)^2+(m-y_4)^2+(m-y_5)^2+(m-y_6)^2 \\ &+(n-z_1)^2+(n-z_2)^2+(n-z_3)^2+(n-z_4)^2+(n-z_5)^2+(n-z_6)^2=C^2. \end{aligned}$$

This equation can be written more briefly thus :

$$\Sigma(l-x_i)^2+\Sigma(m-y_i)^2+\Sigma(n-z_i)^2=C.$$

$$\begin{aligned} \therefore C(l^2+m^2+n^2)-2[l\Sigma(x_i)+m\Sigma(y_i)+n\Sigma(z_i)] \\ +\Sigma(x_i^2)+\Sigma(y_i^2)+\Sigma(z_i^2)=C^2\dots\dots(1). \end{aligned}$$

The equation of the normal is

$$(a^2/x)(l-x)=(b^2/y)(m-y)=(c^2/z)(n-z)\dots\dots(2).$$

$$\text{Also, } x^2/a^2+y^2/b^2+z^2/c^2=1\dots\dots(3).$$

The values of  $y, z$  in terms of  $x$  from (2) in (3) gives

$$x^2/a^2 + \frac{b^2 m^2 x^2}{(a^2 l - a^2 x + b^2 x)^2} + \frac{c^2 n^2 x^2}{(a^2 l - a^2 x + c^2 x)^2} = 1 \dots (4).$$

(4) can be written as follows :

$$x^6 + Ax^5 + Bx^4 + Dx^3 + Ex^2 + Fx + G = 0.$$

Then  $\Sigma(x_1) = -A$ ,  $\Sigma(x_1 x_2) = B$ ,  $\Sigma(x_1^2) = A^2 - 2B$ .

$$\text{But } A = -\frac{2a^2 l}{a^2 - c^2} - \frac{2a^2 l}{a^2 - b^2}.$$

$$B = \frac{a^2(a^2 l^2 + c^2 n^2)}{(a^2 - c^2)^2} + \frac{a^2(a^2 l^2 + b^2 m^2)}{(a^2 - b^2)^2} + \frac{4a^4 l^2}{(a^2 - b^2)(a^2 - c^2)} - a^2.$$

$$\therefore \Sigma(x_1) = \frac{2a^2 l}{a^2 - c^2} + \frac{2a^2 l}{a^2 - b^2}.$$

$$\Sigma(x_1^2) = \frac{2a^2(a^2 l^2 - c^2 n^2)}{(a^2 - c^2)^2} + \frac{2a^2(a^2 l^2 - b^2 m^2)}{(a^2 - b^2)^2} + 2a^2.$$

By symmetry,

$$\Sigma(y_1) = \frac{2b^2 m}{b^2 - c^2} + \frac{2b^2 m}{b^2 - a^2}.$$

$$\Sigma(y_1^2) = \frac{2b^2(b^2 m^2 - c^2 n^2)}{(b^2 - c^2)^2} + \frac{2b^2(b^2 m^2 - a^2 l^2)}{(b^2 - a^2)^2} + 2b^2.$$

$$\Sigma(z_1) = \frac{2c^2 n}{c^2 - b^2} + \frac{2c^2 n}{c^2 - a^2}.$$

$$\Sigma(z_1^2) = \frac{2c^2(c^2 n^2 - b^2 m^2)}{(c^2 - b^2)^2} + \frac{2c^2(c^2 n^2 - a^2 l^2)}{(c^2 - a^2)^2} + 2c^2.$$

Substituting these values in (1) and reducing we get

$$6(l^2 + m^2 + n^2) + \frac{2(c^2 n^2 - a^2 l^2)}{a^2 - c^2} + \frac{2(b^2 m^2 - a^2 l^2)}{a^2 - b^2} + \frac{2(c^2 n^2 - b^2 m^2)}{b^2 - c^2} \\ = C^2 - 2(a^2 + b^2 + c^2).$$

$$\therefore \frac{2(a^4 - 2a^2 b^2 - 2a^2 c^2 + 3b^2 c^2)l^2}{(a^2 - c^2)(a^2 - b^2)(C^2 - 2a^2 - 2b^2 - 2c^2)}$$

$$+ \frac{2(2b^2c^2 + 2a^2b^2 - 3a^2c^2 - b^4)m^2}{(a^2 - b^2)(b^2 - c^2)(C^2 - 2a^2 - 2b^2 - 2c^2)} + \frac{2(c^4 - 2a^2c^2 - 2b^2c^2 + 3a^2b^2)n^2}{(a^2 - c^2)(b^2 - c^2)(C^2 - 2a^2 - 2b^2 - 2c^2)} = 1.$$

$\therefore l^2/R^2 + m^2/S^2 + n^2/T^2 = 1$ , a co-axial ellipsoid.

### CALCULUS.

89. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, O.

Integrate the equation,  $\frac{dy}{dx} + y \cos x = \frac{\sin 2x}{2}$ .

I. Solution by Dr. E. D. ROE, Jr., Norwood, Mass.

Put  $\sin x - y = z$ ; then  $\cos x dx - dy = dz$ ,  $dy = \cos x dx - dz$ , and the equation becomes

$$-\frac{dz}{dx} + (z-1)\cos x = 0, \text{ or } \frac{dz}{z-1} + \cos x dx = 0.$$

Integrating this,  $\log(z-1) + \sin x + \kappa = 0$ ,  $z-1 = e^{-\sin x - \kappa}$ ,

$$\text{or } \sin x - y - 1 = e^{-\sin x - \kappa} = -ce^{-\sin x}, \quad y = \sin x - 1 + ce^{-\sin x}.$$

Dr. Roe also furnished a second solution.

II. Solution by F. ANDEREGG, A. M., Professor of Mathematics, Oberlin College, Oberlin, O.

If the equation  $\frac{dy}{dx} + y \cos x = 0$  is solved, the result is  $y = C_1 e^{-\sin x}$ . After substituting in the original equation  $\frac{dC_1}{dx}$  is found to equal  $e^{\sin x} \sin x \cos x$ ; therefore,  $C_1 = e^{\sin x} \sin x - e^{\sin x} + C$ . And  $y = \sin x + C e^{-\sin x}$ .

III. Solution by WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass., and JOHN R. JEFFERY, Student in Ohio State University, Columbus, O.

The general form of this equation is  $\frac{dy}{dx} + Py = Q$  of which the general integral is  $e^{\int P dx} y = \int e^{\int P dx} Q dx + C$ . Here  $\int P dx = \int \cos x dx = \sin x$ .

$$\therefore e^{\sin x} y = \int e^{\sin x} \cos x \sin x dx + C.$$

Integrating right member by parts,  $e^{\sin x} y = \sin x \cdot e^{\sin x} - e^{\sin x} + c$ , or  $y = \sin x - 1 + ce^{-\sin x}$ .

[See Johnson's *Differential Equations*, page 35, ex. 7.]